arXiv:1508.00362v1 [math.FA] 3 Aug 2015

# EMBEDDINGS INTO ORLICZ SPACES VIA THE MODIFIED RIESZ POTENTIAL

PETTERI HARJULEHTO AND RITVA HURRI-SYRJÄNEN

Abstract. $L^1_1$-functions which are defined in non-smooth domains in the $n$-dimensional Euclidean space can be estimated point-wise by the modified Riesz potential of their gradients. These point-wise estimates imply embeddings into Orlicz spaces from the space $L^1_p$, $1 \le p < n$, where the functions are defined in bounded or unbounded domains with minimum requirement of the smoothness of the boundary. The results are sharp for $L^1_1$-functions.

## 1. Introduction

It is well known that a locally Lipschitz function can be estimated point-wise by the Riesz potential of its gradient in bounded John domains, [20, Theorem], [6, Theorem 10], and hence, especially, in Lipschitz domains and in convex domains, [5, Lemma 7.16]. By modifying the Riesz potential, point-wise estimates can be generalized for functions which are defined in more irregular domains than John domains, [11, Theorem 3.4], [10, Theorem 4.4]. More precisely, for every function $u$ whose weak distributional partial derivatives are in $L^1(G)$, the pointwise estimate

$$|u(x) - u_D| \le \int_G \frac{|\nabla u(y)|}{\psi(|x-y|)^{n-1}}\, dy \tag{1.1}$$

holds for almost every $x \in G$. Here, $G$ is a domain in the $n$-dimensional Euclidean space and the regularity of the boundary is controlled by the function $\psi$. Hedberg's method [13, Lemma, Theorem 1] can be extended so that this point-wise estimate leads to the Sobolev-type inequality where an Orlicz-space is the target space. Hedberg's method has been used by A. Cianchi and B. Stroffolini for the classical Riesz potential when functions are Orlicz functions, [3, Theorem 1, Corollary 1], and by the authors for the modified Riesz potential with a special Orlicz function, [11, Theorem 1.1] and [9, Theorem 1.1], and with a general Orlicz function in [10, Corollary 3.4, Corollary 5.4]. For other papers on Orlicz embeddings of Cianchi we refer to [1], [2].

---

*Date*: August 4, 2015.

2010 *Mathematics Subject Classification.* 31C15 , 42B20, 26D10, 46E30, 46E35.

*Key words and phrases.* Riesz potential, point-wise estimate, Orlicz space, unbounded convex domain, non-smooth domain, Sobolev inequality, Poincaré inequality.

In the present paper we show that the optimal Orlicz function for the modified Riesz potential in (1.1) can be found as a function of $\psi$ which depends on the geometry of the domain $G$. Our main theorem is the following theorem where we give the formula to the Orlicz function.

1.2. **Theorem.** *Let* $1 \le p < n$. *Let the continuous, strictly increasing function* $\varphi : [0,\infty) \to [0,\infty)$ *be such that* $\varphi(0) = \lim_{t\to 0^+} \varphi(t) = 0$ *and* $\varphi$ *satisfies the* $\Delta_2$*-condition and the inequality* $\frac{\varphi(t_1)}{t_1} \le \frac{\varphi(t_2)}{t_2}$ *whenever* $0 < t_1 \le t_2$. *If*

$$\psi(t) = \begin{cases} \varphi(t) & \text{when} \quad 0 \le t \le 1; \\ \varphi(1)t & \text{when} \quad t \ge 1, \end{cases} \tag{1.3}$$

*then there exists an* $N$*-function* $H$ *that satisfies the* $\Delta_2$*-condition, and*

$$H^{-1}(t) \approx \frac{t^{\frac{1}{p}-1}}{\psi\left(t^{-\frac{1}{n}}\right)^{n-1}} \quad \text{for} \quad t > 0.$$

With this function we obtain the following point-wise estimate.

1.4. **Theorem.** *Let* $G$ *be a domain in* $\mathbb{R}^n$, $n \ge 2$. *Let* $1 \le p < n$. *If* $H$ *is the function from Theorem 1.2 and* $\|f\|_{L^p(G)} \le 1$, *then there exists a constant* $C$ *such that the point-wise estimate*

$$H\left(\int_G \frac{|f(y)|}{\psi(|x-y|)^{n-1}}\,dy\right) \le C(Mf(x))^p$$

*holds for every* $x \in \mathbb{R}^n$. *Here,* $Mf$ *is the Hardy-Littelewood maximal operator of* $f$ *and the constant* $C$ *depends on* $n$, $p$, *and the* $\Delta_2$*-constant of* $H$ *only.*

By this point-wise estimate we obtain embedding results for bounded and unbounded non-smooth domains. Examples of these domains are Lipschitz domains and convex domains, but also domains with suitable outward cusps are allowed.

We define a class of domains which are controlled by the function $\psi$ from (1.3). We call these domains in Definition 2.2 as $\varphi$-cigar John domains, since our definition is a modification of [22, 2.1] where J. Väisälä has defined unbounded John domains with $\varphi(t) = t$. Hence, examples of $\varphi$-John domains are the classical bounded and unbounded John domains, but also so called $s$-John domains when $\varphi(t) = t^s$.

We have the following corollary which recovers some of the known results of the Poincaré inequality.

1.5. **Corollary.** *If there exists* $\alpha \in [1, n/(n-1))$ *such that* $t^\alpha/\varphi(t)$ *is increasing for* $t > 0$ *and if* $D$ *is a bounded or an unbounded* $\varphi$*-cigar John domain with a constant* $c_J$ *in* $\mathbb{R}^n$, $n \ge 2$ *and if* $1 \le p < n$, *then with the function* $H$ *in Theorem 1.2 there exists a constant* $C$ *such that the inequality*

$$\inf_{b\in\mathbb{R}} \|u-b\|_{L^H(D)} \le C\|\nabla u\|_{L^p(D)},$$

*holds for every* $u \in L^1_{\text{loc}}(D)$ *with* $|\nabla u| \in L^p(D)$. *Here the constant* $C$ *depends on* $n$, $p$, $\Delta_2$*-constants of* $H$ *and* $\varphi$, *and John constant* $c_J$ *only.*

We point out that if $D$ is a bounded $s$-John domain, then $\varphi(t) = t^s$, $t \geq 0$, and this corollary yields that the $\left(\frac{np}{n-np+sp(n-1)}, p\right)$-Poincaré inequality holds. If $p = 1$, the result is optimal. Thus the corollary recovers some of the known results of [21, Theorem 10], [7, Corollaries 5 and 6], and [17, Theorem 2.3], but our proof is completely different from the previous proofs.

Especially, in Section 6 we construct an example of an unbounded domain which shows that the Lebesque space cannot be the target space in this corresponding embedding if $\lim_{t\to 0^+} t/\varphi(t) = \infty$.

The outline of the paper is as following: We define the domains we consider in Section 2 and we call them $\varphi$-cigar John domains. We find the suitable Orlicz function in Section 3, prove embedding theorems in Section 4, recover some Poincaré inequalities in Section 5, and in Section 6 we construct an example of an unbounded $\varphi$-cigar domain.

## 2. JOHN DOMAINS

Throughout the paper we let the function $\varphi : [0, \infty) \to [0, \infty)$ satisfy the following conditions

(1) $\varphi$ is continuous,
(2) $\varphi$ is strictly increasing,
(3) $\varphi(0) = \lim_{t\to 0^+} \varphi(t) = 0$,
(4) there exists a constant $C_\varphi \geq 1$ such that
$$\frac{\varphi(t_1)}{t_1} \leq C_\varphi \frac{\varphi(t_2)}{t_2}$$
whenever $0 < t_1 \leq t_2$,
(5) $\varphi$ satisfies the $\Delta_2$-condition i.e. there exists a constant $C_\varphi^{\Delta_2} \geq 1$ such that $\varphi(2t) \leq C_\varphi^{\Delta_2}\varphi(t)$ for every $t > 0$.

We write

$$\psi(t) = \begin{cases} \varphi(t) & \text{if} \quad 0 \leq t \leq 1; \\ \varphi(1)t & \text{if} \quad t \geq 1. \end{cases} \tag{2.1}$$

Now, if $\varphi$ satisfies the conditions (1)–(5), then $\psi$ does, too, and the constant in (4) is the same for the functions $\varphi$ and $\psi$, that is $C_\varphi = C_\psi$.

The definition of a bounded John domain goes back to F. John [16, Definition, p. 402] who defined an inner radius and an outer radius domain, and later this domain was renamed as a John domain in [18, 2.1].

We extend the definition of John domains following J. Väisälä [22, 2.1] in the classical case. Let $E$ in $\mathbb{R}^n$, $n \geq 2$, be a closed rectifiable curve with endpoints $a$ and $b$. The subcurve between $x, y \in E$ is denoted by

$E[x,y]$. For $x \in E$ we write

$$q(x) = \min\Big\{\ell\big(E[a,x]\big), \ell\big(E[x,b]\big)\Big\},$$

where $\ell(E[a,x])$ is the length of the subcurve $E[a,x]$.

2.2. **Definition.** A bounded or an unbounded domain $D$ in $\mathbb{R}^n$ is a $\varphi$-cigar John domain if there exists a constant $c_J > 0$ such that each pair of points $a, b \in D$ can be joined by a closed rectifiable curve $E$ in $D$ such that

$$\operatorname{Cig} E(a,b) = \bigcup \left\{ B\left(x, \frac{\psi(q(x))}{c_J}\right) : x \in E \setminus \{a,b\} \right\} \subset D$$

where $B(x,r)$ is an open ball centered at $x$ with a radius $r > 0$ and the function $\psi$ is defined as in (2.1).

The set $\operatorname{Cig} E(a,b)$ is called a cigar with core $E$ joining $a$ and $b$. We point out that if $D$ is a $\varphi$-cigar John domain with $\varphi(t) = t^p$, $p \ge 1$, then it is a $\varphi$-cigar John domain with $\varphi(t) = t^q$ for every $q \ge p$. For the case $\psi(t) = \varphi(t) = t$ for all $t \ge 0$, in Definition 2.2, we refer to [22, 2.1] and [19, 2.11 and 2.13].

If $D$ is a bounded domain then the following definition from [10, Definition 4.1] for a $\psi$-John domain gives an equivalent definition to a bounded $\varphi$-cigar John domain.

2.3. **Definition.** A bounded domain $D$ in $\mathbb{R}^n$, $n \ge 2$, is a $\psi$-John domain if there exist a constants $0 < \alpha \le \beta < \infty$ and a point $x_0 \in D$ such that each point $x \in D$ can be joined to $x_0$ by a rectifiable curve $\gamma : [0, \ell(\gamma)] \to D$, parametrized by its arc length, such that $\gamma(0) = x$, $\gamma(\ell(\gamma)) = x_0$, $\ell(\gamma) \le \beta$, and

$$\psi(t) \le \frac{\alpha}{\ell(\gamma)} \operatorname{dist}(\gamma(t), \partial D) \quad \text{for all} \quad t \in [0, \ell(\gamma)].$$

The point $x_0$ is called a John center of $D$ and $\gamma$ is called a John curve of $x$.

If the function $\psi$ is defined as in (2.1) with the function $\varphi$, then a bounded domain is a $\psi$-John domain if and only if it is a $\varphi$-John domain. If $\psi(t) = t$, then our definition for bounded $\psi$-John domains coincides with the definition of the classical John domains. If $\psi(t) = t^s$, $s \ge 1$, then our definition for bounded $\psi$-John domains coincides with the definition of $s$-John domains.

2.4. **Theorem.** *Let $D$ be a bounded domain. If $D$ is a $\psi$-John domain then $D$ is a $\varphi$-cigar John domain. On the other hand, if $D$ is a $\varphi$-cigar John domain with a constant $c_J$, then $D$ is a $\psi$-John domain with constants*

$$\alpha = \frac{c_J\,\varphi(1)\left(\max\left\{2, \frac{c_J \operatorname{diam}(D)}{\varphi(1)}\right\}\right)^2}{\psi\left(\frac{1}{2c_J}\psi\left(\frac{1}{4}\operatorname{diam}(D)\right)\right)},$$

$$\beta = \max\left\{2, \frac{c_J \operatorname{diam}(D)}{\varphi(1)}\right\}.$$

Note that when $\operatorname{diam}(D) \to \infty$, then $\alpha \to \infty$ with the same speed as $\operatorname{diam}(D)$.

*Proof.* Assume first that $D$ is a $\psi$-John domain with a John center $x_0$. Let $a, b \in D$ and let the John curves $\gamma_1$ and $\gamma_2$ connect them to $x_0$, respectively. We may assume that $a, b \in D \setminus B(x_0, \operatorname{dist}(x_0, \partial D))$, since inside the ball the points can be connect by two straight lines going via the center of the ball $B(x_0, \operatorname{dist}(x_0, \partial D))$. Let $E = \gamma_1 \circ \gamma_2$. Then,

$$\begin{aligned}&\operatorname{Cig} E(a,b)\\ &= \bigcup_{t\in(0,\ell(\gamma_1)]} B\left(\gamma_1(t), \frac{\psi(t)}{\alpha/\operatorname{dist}(x_0,\partial D)}\right) \cup \bigcup_{t\in(0,\ell(\gamma_2)]} B\left(\gamma_2(t), \frac{\psi(t)}{\alpha/\operatorname{dist}(x_0,\partial D)}\right)\end{aligned}$$

and thus $D$ is a $\varphi$-cigar John domain.

Assume then that $D$ is a $\varphi$-cigar John domain. Let us carefully choose a suitable John center so that the center is not too close to the boundary of $D$. Let $x, y \in D$ such that $|x-y| \geq \frac{1}{2}\operatorname{diam}(D)$. Let $E$ be a core of a John cigar that connects$x$ and $y$. Then the length of $E$ is at least $\frac{1}{2}\operatorname{diam}(D)$. Let $x_0$ be the center of $E$. Then

$$\operatorname{dist}(x_0, \partial D) \geq \frac{\psi(\frac{1}{4}\operatorname{diam}(D))}{c_J}$$

so we choose $r = \psi\big(\frac{1}{4}\operatorname{diam}(D)\big)/c_J$, and hence $B(x_0, r) \subset D$. From now on this $r$ and the point $x_0$ are fixed in this proof.

For every $a \in D\setminus\overline{B(x_0,r)}$ there exists a curve $E$ such that $\operatorname{Cig} E(a, x_0) \subset D$. Let $\ell(E)$ be the length of $E$, then $\ell(E) \leq 2$ or by the definition

$$\operatorname{diam}(D) \geq 2\frac{\psi(\ell(E)/2)}{c_J} = 2\frac{\varphi(1)\ell(E)}{2c_J}$$

i.e. $\ell(E) \leq \max\left\{2, \frac{c_J \operatorname{diam}(D)}{\varphi(1)}\right\} = \beta$.



FIGURE 1. The cigar from $a$ to $x_0$ (the solid line), the core $E$ (the dotted line) and a new carrot given by the constant $c_J M$ (the dashed line).

Note that the length of $E$ inside the ball $B(x_0, r)$ is at least $r$ and thus for the points in $E \cap \partial B(x_0, r)$ the distance to the boundary is at least $\psi(r/2)$. Let us choose that

$$M = \frac{\psi(\beta)}{\psi(r/2)} = \frac{\varphi(1)\beta}{\psi(r/2)}.$$

Since $r \le \ell(E) \le \beta$ and $\psi$ is increasing, we have $M \ge 1$.

Let $z_0 \in E$ be the first point from $a$ that satisfies $z_0 \in \partial B(x_0, r)$. Let us replace $E[z_0, x_0]$ by the radius of the ball $B(x_0, r)$, if necessary. Let us denote this new arc by $E$. Let $\gamma$ be an arc $E$ parametrized by its curve length, such that $\gamma(0) = a$, $\gamma(\ell(E)) = x_0$. Since

$$\frac{\psi(\ell(E))}{Mc_J} \le \frac{\psi(r/2)}{c_J}$$

we obtain that

$$\bigcup_{t\in(0,\ell(E))} B\left(\gamma(t), \frac{\psi(t)}{Mc_J}\right) \setminus B(x_0, r) \subset \mathrm{Cig}[a, x_0].$$

This yields that

$$\bigcup_{t\in(0,\ell(E))} B\left(\gamma(t), \frac{\psi(t)}{Mc_J}\right) \subset D$$

and thus

$$\psi(t) \le Mc_J \,\mathrm{dist}(\gamma(t), \partial D) \le \frac{Mc_J\beta}{\ell(E)} \,\mathrm{dist}(\gamma(t), \partial D).$$

This yields that we may choose $\alpha = Mc_J\beta$. Thus, $D$ is a $\psi$-John domain with these $\alpha$ and $\beta$. □

## 3. POINT-WISE ESTIMATES

We note that by the condition (4) of $\varphi$

$$\psi(t) \le C_\varphi \varphi(1)t \quad \text{for all} \quad t \ge 0. \tag{3.1}$$

We recall a covering lemma from [10, 4.3. Lemma] which is valid for a bounded $\varphi$-John domain. For the previous versions in classical case we refer to [8, Theorem 9.3] and in a special case to [11, Lemma 3.5].

3.2. **Lemma.** [10, 4.3. Lemma]. *Let $\varphi$ satisfies the conditions (1)–(5). Let $\psi : [0,\infty) \to [0,\infty)$ be defined as in* (2.1). *Let $D$ in $\mathbb{R}^n$, $n \ge 2$, be a bounded $\psi$-John domain with John constants $\alpha$ and $\beta$. Let $x_0 \in D$ the John center. Then for every $x \in D \setminus B(x_0, \mathrm{dist}(x_0, \partial D))$ there exists a sequence of balls $(B(x_i, r_i))$ such that $B(x_i, 2r_i)$ is in $D$ for each $i = 0, 1, \dots$, and for some constants $K = K(\alpha, \mathrm{dist}(x_0, \partial D), \mathrm{diam}(D), \varphi)$, $N = N(n)$, and $M = M(n)$*

- $B_0 = B\left(x_0, \frac{1}{2}\mathrm{dist}(x_0, \partial D)\right)$;
- $\psi(\mathrm{dist}(x, B_i)) \le Kr_i$, *and* $r_i \to 0$ *as* $i \to \infty$;
- *no point of the domain $D$ belongs to more than $N$ balls $B(x_i, r_i)$; and*

- $|B(x_i,r_i)\cup B(x_{i+1},r_{i+1})| \le M|B(x_i,r_i)\cap B(x_{i+1},r_{i+1})|$.

3.3. *Remark.* (1) The constant $K$ in the previous lemma can be taken to be $K=\max\{2\alpha/\operatorname{dist}(x_0,\partial D), 2\varphi(1), \varphi(\operatorname{diam}(D))/\operatorname{diam}(D)\}$.
(2) If $D$ is a $\varphi$-cigar John domain and the John center has been chosen as in Theorem 2.4, then

$$\frac{\alpha}{\operatorname{dist}(x_0,\partial D)} \le \frac{c_J^2\,\varphi(1)\left(\max\left\{2, \frac{c_J \operatorname{diam}(D)}{\varphi(1)}\right\}\right)^2}{\psi\left(\frac{1}{2c_J}\psi\left(\frac14 \operatorname{diam}(D)\right)\right)\psi\left(\frac14 \operatorname{diam}(D)\right)} \to \frac{32c_J^5}{\varphi(1)^4}$$

as $\operatorname{diam}(D)\to\infty$.

We recall the following definitions. Let $G$ be an open set of $\mathbb{R}^n$. We denote the Lebegue space by $L^p(G)$, $1\le p<\infty$. By $L^1_p(G)$, $1\le p<\infty$, we denote those locally integrable functions whose first weak distributional derivatives belongs to $L^p(G)$ i.e. $L^1_p(G)=\left\{u\in L^1_{\text{loc}}(G) : |\nabla u|\in L^p(G)\right\}$. By $W^{1,p}(G)$, $1\le p<\infty$, we denote those functions from $L^p(G)$ whose first weak distributional derivatives belongs to $L^p(G)$ i.e. $W^{1,p}(G)=\{u\in L^p(G) : |\nabla u|\in L^p(G)\}$.

Theorem 2.4 and Lemma 3.2 give the following point-wise estimate which we recall from [10, 4.4. Theorem].

3.4. **Theorem.** *Let $\varphi$ satisfy the conditions (1)–(5). Let $\psi:[0,\infty)\to[0,\infty)$ be as defined in* (2.1). *Let $D$ in $\mathbb{R}^n, n\ge 2$, be a bounded $\varphi$-cigar John domain with a John constant $c_J$. Then there exists a finite constant $C$ and $x_0\in D$ such that for every $u\in L^1_1(D)$ and for almost every $x\in D$ the inequality*

$$\left|u(x)-u_{B(x_0,\operatorname{dist}(x_0,\partial D))}\right| \le C\int_D \frac{|\nabla u(y)|}{\psi(|x-y|)^{n-1}}\,dy$$

*holds. Here*

$$C=c\left(n, c_J, C_\varphi, C_\varphi^{\Delta_2}, \varphi(1), \min\left\{\operatorname{diam}(D),1\right\}\right).$$

We recall the definitions of $N$-functions and Orlicz spaces.

3.5. **Definition.** A function $H:[0,\infty)\to[0,\infty)$ is an $N$-function if

(N1) $H$ is continuous,
(N2) $H$ is convex,
(N3) $\lim_{t\to0^+}\frac{H(t)}{t}=0$ and $\lim_{t\to\infty}\frac{H(t)}{t}=\infty$.

Continuity and $\lim_{t\to0^+}\frac{H(t)}{t}=0$ yield that $H(0)=0$. Let $0<t<s$ by convexity

$$H(t)=H\left(\frac{t}{s}s+\left(1-\frac{t}{s}\right)0\right)\le\frac{t}{s}H(s)+\left(1-\frac{t}{s}\right)H(0)$$

and thus $\frac{H(t)}{t}\le\frac{H(s)}{s}$ for $0<t<s$.

This implies that $H$ is a strictly increasing function.

By the notation $f \lesssim g$ we mean that there exists a constant $C > 0$ such that $f(x) \leq Cg(x)$ for all $x$. The notation $f \approx g$ means that $f \lesssim g \lesssim f$.

Two $N$-functions $H$ and $K$ are equivalent, which is written as $H \simeq K$, if there exists $m \geq 1$ such that $H(t/m) \leq K(t) \leq H(mt)$ for all $t > 0$. Equivalent $N$-functions give the same space with comparable norms. We point out that $H \simeq K$ if and only if for the inverse functions $H^{-1} \approx K^{-1}$.

We assume that $H$ satisfies the $\Delta_2$-condition, that is, there exists a constant $C_H^{\Delta_2}$ such that

$$H(2t) \leq C_H^{\Delta_2} H(t) \quad \text{for all} \quad t > 0. \tag{3.6}$$

If an $N$-function satisfies the $\Delta_2$-condition then the relations $\simeq$ and $\approx$ are equivalent. The constant $C_H^{\Delta_2}$ is called the $\Delta_2$-constant of $H$.

Let $G$ in $\mathbb{R}^n$ be an open set. The Orlicz class is a set of all measurable functions $u$ defined on $G$ such that

$$\int_G H\big(|u(x)|\big)\, dx < \infty\,.$$

We study the Orlicz space $L^H(G)$ which means the space of all measurable functions $u$ defined on $G$ such that

$$\int_G H\big(\lambda |u(x)|\big)\, dx < \infty$$

for some $\lambda > 0$.

Whenever the function $H$ satisfies the $\Delta_2$-condition, then the space $L^H(G)$ is a vector space and it is equivalent to the corresponding Orlicz class. We study these Orlicz spaces and call their functions Orlicz functions. The Orlicz space $L^H(G)$ equipped with the Luxemburg norm

$$||u||_{L^\Phi(G)} = \inf\left\{\lambda > 0 : \int_G \Phi\left(\frac{|u(x)|}{\lambda}\right) dx \leq 1\right\}$$

is a Banach space.

We recall the following theorem from [10, 1.3. Theorem].

3.7. **Theorem.** *Let $\varphi$ satisfy the conditions (1)–(5). Let $\psi : [0, \infty) \to [0, \infty)$ be defined as in* (2.1). *Let $1 \leq p < n$ be given. Suppose that there exists a continuous function $h : [0, \infty) \to [0, \infty)$ such that*

$$\sum_{k=1}^{\infty} \frac{(2^{-k}t)^n}{\psi(t2^{-k})^{n-1}} \leq h(t) \quad \textit{for all} \quad t > 0\,. \tag{3.8}$$

*Let $\delta : (0, \infty) \to [0, \infty)$ be a continuous function and let $H : [0, \infty) \to [0, \infty)$ be an N-function satisfying the $\Delta_2$-condition. Suppose that there exists a finite constant $C_H$ such that the inequality*

$$H\left(h(\delta(t))t + \psi(\delta(t))^{1-n}(\delta(t))^{n(1-\frac{1}{p})}\right) \leq C_H t^p \tag{3.9}$$

*holds for all* $t > 0$. *Let* $G$ *in* $\mathbb{R}^n$ *be an open set. If* $\|f\|_{L^p(G)} \leq 1$, *then there exists a constant* $C$ *such that the inequality*

$$H\left(\int_G \frac{|f(y)|}{\psi(|x-y|)^{n-1}}\,dy\right) \leq C(Mf(x))^p \tag{3.10}$$

*holds for every* $x \in \mathbb{R}^n$. *Here the constant* $C$ *depends on* $n$, $p$, $C_\varphi$, $C_H$, *and the* $\Delta_2$*-constants of* $\varphi$ *and* $H$ *only.*

Our goal is to find a formula which would give all suitable functions $H$. Examples of some of these functions were given in [10, Section 6].

Here we do the preparations to find $H$. Assume that there exists $\alpha \in [1, n/(n-1))$ such that $t^\alpha/\varphi(t)$ is increasing for $t > 0$. This yields that $t^\alpha/\psi(t)$ is increasing, too. Under this condition inequality (3.8) holds: Since

$$\begin{aligned}\frac{(2^{-k}t)^n}{\psi(t2^{-k})^{n-1}} &= \frac{(2^{-k}t)^n}{(2^{-k}t)^{\alpha(n-1)}}\cdot\frac{(2^{-k}t)^{\alpha(n-1)}}{\psi(t2^{-k})^{n-1}}\\ &\leq (2^{-k}t)^{n-\alpha(n-1)}\frac{t^{\alpha(n-1)}}{\psi(t)^{n-1}} = 2^{-k(n-\alpha(n-1))}\frac{t^n}{\psi(t)^{n-1}},\end{aligned}$$

we have

$$\sum_{k=1}^{\infty}\frac{(2^{-k}t)^n}{\psi(t2^{-k})^{n-1}} \leq C(n,\alpha)\frac{t^n}{\psi(t)^{n-1}}, \quad \text{where} \quad C(n,\alpha) = \frac{2^{\alpha(n-1)}}{2^n - 2^{\alpha(n-1)}}.$$

Let us define the functions $h$ and $\delta$ such that

$$h(t) = C(n,\alpha)\frac{t^n}{\psi(t)^{n-1}} \quad \text{and} \quad \delta(t) = t^{-\frac{p}{n}} \quad \text{for all} \quad t > 0.$$

Then,

$$\begin{aligned}h(\delta(t))t + \psi(\delta(t))^{1-n}(\delta(t))^{n(1-\frac{1}{p})} &= h\left(t^{-\frac{p}{n}}\right)t + \psi\left(t^{-\frac{p}{n}}\right)^{1-n}\left(t^{-\frac{p}{n}}\right)^{n(1-\frac{1}{p})}\\ &= \frac{C(n,\alpha)t^{-p}}{\psi\left(t^{-\frac{p}{n}}\right)^{n-1}}t + \frac{t^{1-p}}{\psi\left(t^{-\frac{p}{n}}\right)^{n-1}}\\ &= \frac{(C(n,\alpha)+1)t^{1-p}}{\psi\left(t^{-\frac{p}{n}}\right)^{n-1}}.\end{aligned}$$

If we choose

$$F^{-1}(t) = \frac{(C(n,\alpha)+1)(t^{1/p})^{1-p}}{\psi\left((t^{1/p})^{-\frac{p}{n}}\right)^{n-1}} = \frac{(C(n,\alpha)+1)t^{\frac{1}{p}-1}}{\psi\left(t^{-\frac{1}{n}}\right)^{n-1}}$$

and assume that the inverse function of $F^{-1}$ exists, that is $(F^{-1})^{-1} =: F$ exists, then

$$h(\delta(t))t + \psi(\delta(t))^{1-n}(\delta(t))^{n(1-\frac{1}{p})} = F^{-1}(t^p)$$

and thus

$$F\left(h(\delta(t))t + \psi(\delta(t))^{1-n}(\delta(t))^{n(1-\frac{1}{p})}\right) = F\left(F^{-1}(t^p)\right) = t^p.$$

Unfortunately, there is a problem with this function $F$ to be a suitable function $H$; namely, the function $F$ is not necessary convex. For example, if $n = 2$, $\varphi(t) = t^{\frac{3}{2}}$, and $p = 1.9$, then the function $F$ is not convex, see Figure 2. The angle at the point $(1, F^{-1}(1))$ comes from the angle of $\psi$ at the point $(1, \psi(1))$. Our main theorem, Theorem 1.2 in Introduction, corrects this point: we show that there exists an $N$-function $H$ that is equivalent with $F$.



FIGURE 2. The function $F$ is not necessary convex.

*Proof of Theorem 1.2.* Let us write that

$$F^{-1}(t) = \frac{t^{\frac{1}{p}-1}}{\psi\left(t^{-\frac{1}{n}}\right)^{n-1}}$$

for $t > 0$ and $F^{-1}(0) = 0$. Let us first show that $F^{-1}$ is strictly increasing. Assume then that $0 < s < t$. The inequality $F^{-1}(s) < F^{-1}(t)$ is equivalent to the inequality

$$\frac{\psi\left(\left(\frac{1}{t}\right)^{\frac{1}{n}}\right)^{n-1}}{\left(\frac{1}{t}\right)^{1-\frac{1}{p}}} < \frac{\psi\left(\left(\frac{1}{s}\right)^{\frac{1}{n}}\right)^{n-1}}{\left(\frac{1}{s}\right)^{1-\frac{1}{p}}}.$$

Recall that if $\varphi$ satisfies the condition (4), then $\psi$ does, too, and the constant is the same for both functions. Thus by the condition (4) and the inequality $p < n$ we obtain

$$\frac{\psi\left(\left(\frac{1}{t}\right)^{\frac{1}{n}}\right)^{n-1}}{\left(\frac{1}{t}\right)^{1-\frac{1}{p}}} = \left(\frac{\psi\left(\left(\frac{1}{t}\right)^{\frac{1}{n}}\right)}{\left(\frac{1}{t}\right)^{\frac{1}{n}}}\right)^{n-1} \left(\frac{1}{t}\right)^{\frac{n-1}{n}-1+\frac{1}{p}} \leq \left(\frac{\psi\left(\left(\frac{1}{s}\right)^{\frac{1}{n}}\right)}{\left(\frac{1}{s}\right)^{\frac{1}{n}}}\right)^{n-1} \left(\frac{1}{t}\right)^{\frac{n-1}{n}-1+\frac{1}{p}}$$

$$< \left(\frac{\psi\left(\left(\frac{1}{s}\right)^{\frac{1}{n}}\right)}{\left(\frac{1}{s}\right)^{\frac{1}{n}}}\right)^{n-1} \left(\frac{1}{s}\right)^{\frac{n-1}{n}-1+\frac{1}{p}} = \frac{\psi\left(\left(\frac{1}{s}\right)^{\frac{1}{n}}\right)^{n-1}}{\left(\frac{1}{s}\right)^{1-\frac{1}{p}}}.$$

Thus the function $F^{-1}$ is strictly increasing. This yields that the function $F$ exists and is strictly increasing.

Let us show that $\lim_{t\to 0^+} F^{-1}(t) = 0$. Since $p < n$ we obtain

$$\lim_{t\to 0^+} F^{-1}(t) = \lim_{t\to 0^+} \frac{t^{\frac{1}{p}-1}}{\psi\left(t^{-\frac{1}{n}}\right)^{n-1}} = \lim_{t\to 0^+} \varphi(1)^{1-n} t^{\frac{n-1}{n}+\frac{1}{p}-1} = 0.$$

Let us show that $\lim_{t\to\infty} F^{-1}(t) = \infty$. Since $t/\varphi(t)$ is decreasing, by the condition (4), and by $p < n$ we obtain

$$\lim_{t\to\infty} F^{-1}(t) = \lim_{t\to\infty} \frac{t^{\frac{1}{p}-1}}{\psi\left(t^{-\frac{1}{n}}\right)^{n-1}} = \lim_{t\to\infty} t^{\frac{1}{p}-\frac{1}{n}} \left(\frac{t^{-\frac{1}{n}}}{\psi\left(t^{-\frac{1}{n}}\right)}\right)^{n-1} \geq \lim_{t\to\infty} \frac{t^{\frac{1}{p}-\frac{1}{n}}}{\varphi(1)^{n-1}} = \infty.$$

We have shown that $F^{-1} : [0,\infty) \to [0,\infty)$ is bijective.

Let us then study the condition

$$\frac{F(s)}{s} < \frac{F(t)}{t} \quad \text{for} \quad 0 < s < t. \tag{3.11}$$

Since $F^{-1}$ is a strictly increasing bijection, inequality (3.11) is equivalent to

$$\frac{s}{F^{-1}(s)} < \frac{t}{F^{-1}(t)}.$$

Since $t^\alpha/\varphi(t)$ is increasing, then $\varphi(t)/t^\alpha$ is decreasing and $\psi(t)/t^\alpha$ is decreasing, too. We note that $1 - \frac{\alpha(n-1)}{n} > 0$, since $\alpha < \frac{n}{n-1}$. We obtain

$$\begin{aligned}
\frac{s}{F^{-1}(s)} &= s^{2-\frac{1}{p}}\psi\left(s^{-\frac{1}{n}}\right)^{n-1} = s^{2-\frac{1}{p}-\frac{\alpha(n-1)}{n}}\left(\frac{\psi\left(s^{-\frac{1}{n}}\right)}{\left(s^{-\frac{1}{n}}\right)^\alpha}\right)^{n-1}\\
&= s^{1-\frac{1}{p}+1-\frac{\alpha(n-1)}{n}}\left(\frac{\psi\left(s^{-\frac{1}{n}}\right)}{\left(s^{-\frac{1}{n}}\right)^\alpha}\right)^{n-1} < t^{1-\frac{1}{p}+1-\frac{\alpha(n-1)}{n}}\left(\frac{\psi\left(t^{-\frac{1}{n}}\right)}{\left(t^{-\frac{1}{n}}\right)^\alpha}\right)^{n-1} = \frac{t}{F^{-1}(t)}
\end{aligned}$$

and thus inequality (3.11) holds.

Let us then show that $F^{-1}(cs) \geq 2F^{-1}(s)$ for all $s \geq 0$ with $c = 2^{\frac{np}{n-p}}$. The inequality $F^{-1}(cs) \geq 2F^{-1}(s)$ is equivalent to

$$2\frac{\psi\left(\left(\frac{1}{cs}\right)^{\frac{1}{n}}\right)^{n-1}}{\left(\frac{1}{cs}\right)^{1-\frac{1}{p}}} \leq \frac{\psi\left(\left(\frac{1}{s}\right)^{\frac{1}{n}}\right)^{n-1}}{\left(\frac{1}{s}\right)^{1-\frac{1}{p}}}.$$

By the condition $(4)$ of $\varphi$ and the inequality $p < n$, we obtain

$$2\frac{\psi\left(\left(\frac{1}{cs}\right)^{\frac{1}{n}}\right)^{n-1}}{\left(\frac{1}{cs}\right)^{1-\frac{1}{p}}} = 2\left(\frac{\psi\left(\left(\frac{1}{cs}\right)^{\frac{1}{n}}\right)}{\left(\frac{1}{cs}\right)^{\frac{1}{n}}}\right)^{n-1}\left(\frac{1}{cs}\right)^{\frac{n-1}{n}-1+\frac{1}{p}} = \left(\frac{\psi\left(\left(\frac{1}{cs}\right)^{\frac{1}{n}}\right)}{\left(\frac{1}{cs}\right)^{\frac{1}{n}}}\right)^{n-1}\left(\frac{1}{s}\right)^{\frac{n-1}{n}-1+\frac{1}{p}}$$
$$\leq \left(\frac{\psi\left(\left(\frac{1}{s}\right)^{\frac{1}{n}}\right)}{\left(\frac{1}{s}\right)^{\frac{1}{n}}}\right)^{n-1}\left(\frac{1}{s}\right)^{\frac{n-1}{n}-1+\frac{1}{p}} = \frac{\psi\left(\left(\frac{1}{s}\right)^{\frac{1}{n}}\right)^{n-1}}{\left(\frac{1}{s}\right)^{1-\frac{1}{p}}}.$$

The inequality $F^{-1}(cs) \geq 2F^{-1}(s)$ yields that $F$ satisfies the $\Delta_2$-condition. Let us write $F(t) = s$. Then $F^{-1}(s) = t$. Since $F$ is increasing, we have

$$F(2t) = F(2F^{-1}(s)) \leq F(F^{-1}(cs)) = cs = cF(t).$$

P. Hästö has shown in [15, Proposition 5.1] that if $f : [0,\infty) \to [0,\infty)$ satisfies the $\Delta_2$-condition and $x \mapsto f(x)/x$ is increasing, then $f$ is equivalent to a convex function. Since $F$ satisfies inequality (3.11) and the $\Delta_2$-condition, we obtain that $F$ is equivalent to a convex function $H$.

Using $\lim_{t\to 0^+} F^{-1}(t) = 0$ and the bijectivity, we obtain

$$\lim_{t\to 0^+}\frac{F(t)}{t} = \lim_{t\to 0^+}\frac{t}{F^{-1}(t)} = \lim_{t\to 0^+}\frac{t\,\psi\left(\left(\frac{1}{t}\right)^{\frac{1}{n}}\right)^{n-1}}{\left(\frac{1}{t}\right)^{1-\frac{1}{p}}} = \lim_{t\to 0^+}\varphi(1)^{n-1}t^{1-\frac{1}{p}+1-\frac{n-1}{n}} = 0$$

and thus also $\lim_{t\to 0^+}\frac{H(t)}{t} = 0$. This gives that $H$ is right continuous at the origin. Thus by convexity the function $H$ is continuous on $[0,\infty)$.

Since $\varphi(t)/t^\alpha$ is decreasing and $\alpha < \frac{n}{n-1}$, we obtain

$$\lim_{t\to\infty}\frac{F(t)}{t} = \lim_{t\to\infty}\frac{t}{F^{-1}(t)} = \lim_{t\to\infty} t^{2-\frac{1}{p}}\varphi\left(t^{-\frac{1}{n}}\right)^{n-1}$$
$$= \lim_{t\to\infty} t^{2-\frac{1}{p}-\frac{\alpha(n-1)}{n}}\left(\frac{\varphi\left(t^{-\frac{1}{n}}\right)}{\left(t^{-\frac{1}{n}}\right)^{\alpha}}\right)^{n-1} \geq \lim_{t\to\infty} t^{1-\frac{1}{p}+1-\frac{\alpha(n-1)}{n}}\left(\frac{\varphi(1)}{1^\alpha}\right)^{n-1} = \infty.$$

Since the functions $F$ and $H$ are equivalent, this yields that

$$\lim_{t\to\infty}\frac{H(t)}{t} = \infty.$$

Thus we have shown that the function $H$ satisfies the conditions (N1) – (N3). □

3.12. *Remark.* Later it is crucial to us that

$$H^{-1}(t) \approx \frac{t^{\frac{1}{p}-1}}{\psi\left(t^{-\frac{1}{n}}\right)^{n-1}} = \frac{t^{\frac{1}{p}-1}}{\varphi(1)^{n-1}\left(t^{-\frac{1}{n}}\right)^{n-1}} = \varphi(1)^{1-n}t^{\frac{n-p}{np}}$$

for $0 < t \leq 1$. Namely, then for every $\varphi$ the function $H$ satisfies $H(t) \approx t^{\frac{np}{n-p}}$ whenever $0 < t \leq 1$.

3.13. **Example.** Functions $\varphi(t) = t^\alpha / \log^\beta(e+1/t)$, $\alpha \in [1, \frac{n}{n-1})$ and $\beta \geq 0$, satisfy the assumptions of Theorem 1.2.

Now, the proof for our second main theorem, Theorem 1.4 in Introduction, follows easily:

*Proof of Theorem 1.4.* Theorem 3.7 and Theorem 1.2. □

As a corollary we obtain from Theorem 1.4 and Theorem 3.4:

3.14. **Corollary.** *Let* $1 \leq p < n$. *Let the function* $H$ *be as in Theorem 1.2. If* $D$ *is a bounded* $\varphi$*-cigar John domain with a constant* $c_J$, *then there exit a constant* $C$ *and a point* $x_0 \in D$ *such that the point-wise estimate*

$$H\left(\left|u(x) - u_{B(x_0,\operatorname{dist}(x_0,\partial D))}\right|\right) \leq C(M|\nabla u|(x))^p$$

*holds for all* $u \in L^1_p(D)$ *with* $\|\nabla u\|_{L^p(D)} \leq 1$ *and for almost every* $x \in D$. *Here the constant* $C$ *depends on* $n$, $p$, $C_H$, $C_H^{\Delta_2}$, $C_\varphi^{\Delta_2}$, $c_J$, $\varphi(1)$ *and* $\min\left\{\operatorname{diam}(D), 1\right\}$ *only.*

## 4. On embeddings

Corollary 3.14 is essential in the proofs of the following Theorem 4.1 and Theorem 4.3.

4.1. **Theorem** (Bounded domain, $1 < p < n$). *Assume that* $\varphi$ *satisfies the conditions* (1)– (5), $C_\varphi = 1$ *in the condition* (4), *and there exists* $\alpha \in [1, n/(n-1))$ *such that* $t^\alpha/\varphi(t)$ *is increasing for* $t > 0$. *Let* $\psi$ *be defined as in* (2.1). *Let* $D \subset \mathbb{R}^n$, $n \geq 2$, *be a bounded* $\varphi$*-cigar John domain with a constant* $c_J$. *Let* $1 < p < n$. *Then there exists an* $N$*-function* $H$, *that satisfies* $\Delta_2$*-condition and*

$$H^{-1}(t) \approx \frac{t^{\frac{1}{p}-1}}{\psi\left(t^{-\frac{1}{n}}\right)^{n-1}} \quad \textit{for all } t > 0\,,$$

*and there exists a constant* $C < \infty$ *such that the inequality*

$$\|u - u_D\|_{L^H(D)} \leq C\|\nabla u\|_{L^p(D)},$$

*holds for every* $u \in L^1_p(D)$. *Here the constant* $C$ *depends on* $n$, $p$, $C_H^{\Delta_2}$, $C_\varphi^{\Delta_2}$, $c_J$ *and* $\min\{\operatorname{diam}(D), 1\}$ *only.*

*Proof.* Assume that $\|\nabla u\|_{L^p(D)} \leq 1$. Corollary 3.14 yields that

$$H\left(\left|u(x) - u_{B(x_0,\operatorname{dist}(x_0,\partial D))}\right|\right) \leq C(M|\nabla u|(x))^p,$$

where the constant $C$ depends on $n$, $p$, $C_H^{\Delta_2}$, $C_\varphi^{\Delta_2}$, $c_J$, and $\min\{\mathrm{diam}(D), 1\}$ only. By integrating over $D$ and using the fact that the maximal operator is bounded whenever $1 < p < n$, we obtain that

$$\begin{aligned}\int_D H\left(\left|u(x) - u_{B(x_0,\mathrm{dist}(x_0,\partial D))}\right|\right)\,dx &\le C\int_D (M|\nabla u|(x))^p\,dx\\ &\le C\int_D |\nabla u(x)|^p\,dx \le C.\end{aligned}$$

This yields that the inequality

$$\|u - u_{B(x_0,\mathrm{dist}(x_0,\partial D))}\|_{L^H(D)} \le C$$

holds for every $u \in L^1_p(D)$ with $\|\nabla u\|_{L^p(D)} \le 1$. By applying this inequality to the function $u/\|\nabla u\|_{L^p(D)}$ we obtain that

$$\|u - u_{B(x_0,\mathrm{dist}(x_0,\partial D))}\|_{L^H(D)} \le C\|\nabla u\|_{L^p(D)}.$$

We may assume w.l.o.g. that $\|\nabla u\|_{L^p(D)} \ne 0$. Let denote $B = B(x_0, \mathrm{dist}(x_0, \partial D))$. By the triangle inequality

$$\|u - u_D\|_{L^H(D)} \le \|u - u_B\|_{L^H(D)} + \|u_B - u_D\|_{L^H(D)}.$$

Here,

$$\begin{aligned}\|u_B - u_D\|_{L^H(D)} = |u_B - u_D|\,\|1\|_{L^H(D)} &\le \frac{\|1\|_{L^H(D)}}{|D|}\|u - u_B\|_{L^1(D)}\\ &\le C\frac{\|1\|_{L^H(D)}\|1\|_{L^{H^*}(D)}}{|D|}\|u - u_B\|_{L^H(D)}\end{aligned}$$

where $H^*$ is the conjugate function of $H$ and $C$ is the constant in Hölder's inequality.

Next we show that $\|1\|_{L^H(D)}\|1\|_{L^{H^*}(D)} \approx |D|$. Since the function $H$ is continuous and strictly increasing, there exists a unique $\lambda > 0$ such that

$$H(1/\lambda)\,|D| = \int_D H(1/\lambda)\,dx = 1$$

i.e. $\lambda = \|1\|_{L^H(D)}$. By solving $\lambda$ we obtain

$$\|1\|_{L^H(D)} = \frac{1}{H^{-1}\left(\frac{1}{|D|}\right)}.$$

Similarly, we obtain

$$\|1\|_{L^{H^*}(D)} = \frac{1}{(H^*)^{-1}\left(\frac{1}{|D|}\right)}.$$

Since

$$t \le H^{-1}(t)(H^*)^{-1}(t) \le 2t$$

for all $t \ge 0$, see for example [4, Lemma 2.6, p. 56], we obtain that

$$\|1\|_{L^H(D)}\|1\|_{L^{H^*}(D)} = \frac{1}{H^{-1}\left(\frac{1}{|D|}\right)(H^*)^{-1}\left(\frac{1}{|D|}\right)} \le |D|.$$

Hence, we have shown that

$$\|u - u_D\|_{L^H(D)} \le C\|\nabla u\|_{L^p(D)}$$

for every $u \in L^1_p(D)$. □

4.2. **Example.** Let us choose that $\varphi(t) = t^s$, $s \in [1, \frac{n}{n-1})$. We have calculated in Remark 3.12 that for every $\varphi$ the function $H$ satisfies $H(t) \approx t^{\frac{np}{n-p}}$ whenever $0 < t \le 1$. If $t > 1$, then

$$H^{-1}(t) \approx \frac{t^{\frac{1}{p}-1}}{\psi\left(t^{-\frac{1}{n}}\right)^{n-1}} = \frac{t^{\frac{1}{p}-1}}{\varphi\left(t^{-\frac{1}{n}}\right)^{n-1}} = \frac{t^{\frac{1}{p}-1}}{\left(t^{-\frac{1}{n}}\right)^{s(n-1)}} = t^{\frac{n-np+sp(n-1)}{np}}$$

and thus we have that $H(t) \approx t^{\frac{np}{n-np+sp(n-1)}}$ for $t > 1$.

4.3. **Theorem** (Bounded domain, $p = 1$)**.** *Assume that the function $\varphi$ satisfies the conditions (1)– (5), $C_\varphi = 1$ in the condition (4), and there exists $\alpha \in [1, n/(n-1))$ such that $t^\alpha/\varphi(t)$ is increasing for $t > 0$. Let $\psi$ be defined as in (2.1) Let $D \subset \mathbb{R}^n$, $n \ge 2$, be a bounded $\varphi$-cigar John domain with a constant $c_J$. Then there exists an N-function $H$, that satisfies $\Delta_2$-condition and*

$$H^{-1}(t) \approx \frac{1}{\psi\left(t^{-\frac{1}{n}}\right)^{n-1}} \quad \textit{for all } t > 0\,,$$

*such that the inequality*

$$\|u - u_D\|_{L^H(D)} \le C\|\nabla u\|_{L^1(D)},$$

*holds for some constant $C$ and for every $u \in L^1_p(D)$. Here the constant $C$ depends on $n$, $C_H^{\Delta_2}$, $C_\varphi^{\Delta_2}$ and $c_J$ only.*

*Proof.* Let us consider functions $u \in L^1_1(D)$ such that $\|\nabla u\|_{L^1(D)} \le 1$. The center ball $B(x_0, \operatorname{dist}(x_0, \partial D))$ is written as $B$. In the proof of Theorem 2.4 we had chosen $x_0$ so that $\operatorname{dist}(x_0, \partial D) \ge \psi(\frac{1}{4}\operatorname{diam}(D))/c_J$. We show that there exists a constant $C < \infty$ such that the inequality

$$\int_D H(|u(x) - u_B|)\,dx \le C \tag{4.4}$$

holds whenever $\|\nabla u\|_{L^1(D)} \le 1$. This yields the claim as in the proof of Theorem 4.1.

Since $H$ is increasing, we first estimate

$$\int_D H(|u(x) - u_B|)\,dx \le \sum_{j \in \mathbb{Z}} \int_{\{x \in D : 2^j < |u(x) - u_B| \le 2^{j+1}\}} H(2^{j+1})\,dx.$$

Let us define

$$v_j(x) = \max\left\{0, \min\left\{|u(x) - u_B| - 2^j, 2^j\right\}\right\}$$

for all $x \in D$. If $x \in \{x \in D : 2^j < |u(x) - u_B| \le 2^{j+1}\}$, then $v_{j-1}(x) \ge 2^{j-1}$. We obtain

$$\int_D H(|u(x) - u_B|)\,dx \le \sum_{j\in\mathbb{Z}} \int_{\{x\in D: v_j(x)\ge 2^j\}} H(2^{j+2})\,dx. \tag{4.5}$$

By the triangle inequality we have

$$v_j(x) = |v_j(x) - (v_j)_B + (v_j)_B| \le |v_j(x) - (v_j)_B| + |(v_j)_B|.$$

By the $(1,1)$-Poincaré inequality in a ball $B$, [5, Section 7.8], there exists a constant $C(n)$ such that

$$\begin{aligned} |(v_j)_B| = (v_j)_B &= \fint_B v_j(x)\,dx \le \fint_B |u(x) - u_B|\,dx \\ &\le C(n)|B|^{\frac{1}{n}} \fint_B |\nabla u(x)|\,dx \le C(n)|B|^{\frac{1}{n}-1}. \end{aligned}$$

We continue to estimate the right hand side of inequality (4.5)

$$\begin{aligned} \int_D H(|u(x) - u_B|)\,dx &\le \sum_{j\in\mathbb{Z}} \int_{\{x\in D: |v_j(x)-(v_j)_B| + C|B|^{-1}\ge 2^j\}} H(2^{j+2})\,dx \\ &\le \sum_{j\in\mathbb{Z}} \int_{\{x\in D: |v_j(x)-(v_j)_B|\ge 2^{j-1}\}} H(2^{j+2})\,dx + \sum_{2^{j-1}\le C(n)|B|^{\frac{1}{n}-1}} \int_D H(2^{j+2})\,dx \\ &\le \sum_{j\in\mathbb{Z}} \int_{\{x\in D: |v_j(x)-(v_j)_B|\ge 2^{j-1}\}} H(2^{j+2})\,dx + \sum_{j=-\infty}^{j_0} \int_D H(2^{j+2})\,dx, \end{aligned} \tag{4.6}$$

where $j_0 = \lceil \log(C(n)|B|^{\frac{1}{n}-1}) \rceil$.

Assume first that $\operatorname{diam}(D)$ is so large that $j_0 \le -2$. When $t < 1$, then $\psi(t^{-1/n}) = \varphi(1)t^{-1/n}$ by (2.1) and thus

$$H^{-1}(t) = \frac{1}{\psi(t^{-1/n})^{n-1}} = \varphi(1)^{1-n} t^{\frac{n-1}{n}}.$$

Thus for $t < 1$ we obtain that $H(t) \approx t^{\frac{n}{n-1}}$. This yields that

$$\begin{aligned} \sum_{j=-\infty}^{j_0} \int_D H(2^{j+2})\,dx &\approx |D| \sum_{j=-\infty}^{\lceil \log(C|B|^{\frac{1}{n}-1})\rceil} 2^{\frac{n(j+2)}{n-1}} \le C|D| 2^{\frac{n}{n-1}\cdot\lceil \log(C|B|^{\frac{1}{n}-1})\rceil} \\ &\le C|D||B|^{\frac{n}{n-1}(\frac{1}{n}-1)} = C|D||B|^{-1} \\ &\le C \frac{\operatorname{diam}(D)^n}{(\psi(\frac{1}{4}\operatorname{diam}(D))/c_J)^n}. \end{aligned} \tag{4.7}$$

This constant does not blow up when $\operatorname{diam}(D) \to \infty$:

$$\frac{\operatorname{diam}(D)^n}{(\psi(\frac{1}{4}\operatorname{diam}(D))/c_J)^n} \to \frac{4^n c_J^n}{\varphi(1)^n} \quad \text{as} \quad \operatorname{diam}(D) \to \infty.$$

Assume then that $\operatorname{diam}(D)$ is small. This yields that for every $j_0 \in \mathbb{Z}$ the sum $\sum_{j=-2}^{j_0} H(2^{j+2})$ is finite and depends on $j_0$. We obtain

$$\sum_{j=-\infty}^{j_0} \int_D H(2^{j+2})\,dx \le \sum_{j=-\infty}^{-2} \int_D H(2^{j+2}) + \sum_{j=-2}^{j_0} H(2^{j+2}) < \infty. \tag{4.8}$$

Then, we will find an upper bound for the sum

$$\sum_{j\in\mathbb{Z}} \int_{\{x\in D:|v_j(x)-(v_j)_B|\ge 2^{j-1}\}} H(2^{j+2})\,dx.$$

Since $\|\nabla v_j\|_{L^1(D)} \le \|\nabla u\|_{L^1(D)} \le 1$, Corollary 3.14 yields that

$$\begin{aligned}\sum_{j\in\mathbb{Z}} \int_{\{x\in D:|v_j(x)-(v_j)_B|\ge 2^{j-1}\}} H(2^{j+2})\,dx &= \sum_{j\in\mathbb{Z}} \int_{\{x\in D:H(|v_j(x)-(v_j)_B|)\ge H(2^{j-1})\}} H(2^{j+2})\,dx\\ &\le \sum_{j\in\mathbb{Z}} \int_{\{x\in D:CM|\nabla v_j|(x)\ge H(2^{j-1})\}} H(2^{j+2})\,dx.\end{aligned}$$

We choose for every $x \in \{x \in D : CM|\nabla v_j|(x) \ge H(2^{j-2})\}$ a ball $B(x, r_x)$, centered at $x$ and with radius $r_x$ depending on $x$, such that

$$C \fint_{B(x,r_x)} |\nabla v_j(y)|\,dy \ge \frac{1}{2} H(2^{j-1})$$

with the understanding that $|\nabla v_j|$ is zero outside $D$. By the Besicovitch covering theorem (or the 5-covering theorem) we obtain a subcovering $\{B_k\}_{k=1}^{\infty}$ so that we may estimate by the $\Delta_2$-condition of $H$

$$\begin{aligned}\sum_{j\in\mathbb{Z}} \int_{\{x\in D:|v_j(x)-(v_j)_B|\ge 2^{j-1}\}} H(2^{j+2})\,dx &\le \sum_{j\in\mathbb{Z}} \sum_{k=1}^{\infty} \int_{B_k} H(2^{j+2})\,dx\\ \le \sum_{j\in\mathbb{Z}} \sum_{k=1}^{\infty} |B_k| H(2^{j+2}) &\le \sum_{j\in\mathbb{Z}} \sum_{k=1}^{\infty} C|B_k| \frac{H(2^{j+2})}{H(2^{j-1})} \fint_{B_k} |\nabla v_j(y)|\,dy\\ \le C \sum_{j\in\mathbb{Z}} \int_D |\nabla v_j(y)|\,dy.\end{aligned}$$

Let $E_j = \{x \in D : 2^j < |u(x) - u_B| \le 2^{j+1}\}$. Since $|\nabla v_j|$ is zero almost everywhere in $D \setminus E_j$ and $|\nabla u(x)| = \sum_j |\nabla v_j(x)|\chi_{E_j}(x)$ for almost every $x \in D$, we obtain

$$\sum_{j\in\mathbb{Z}} \int_{\{x\in D:|v_j(x)-(v_j)_B|\ge 2^{j-1}\}} H(2^{j+2})\,dx \le C \int_D |\nabla u(y)|\,dy \le C. \tag{4.9}$$

Estimates (4.6), (4.7), (4.8) and (4.9) imply inequality (4.4). □

4.10. *Remark.* Corollary 1.5 in Introduction follows from Theorem 4.1 and Theorem 4.3.

4.11. *Remark.* In Theorem 4.3 the $N$-function $H$ is the best possible in a sense that it cannot be replaced by any $N$-function $K$ that satisfies the $\Delta_2$-condition and

$$\lim_{t\to\infty}\frac{K(t)}{H(t)}=\infty.$$

In [10, Theorem 7.2] we have shown that the corresponding embedding in Theorem 4.3 does not hold if

$$\lim_{t\to 0^+} t^n K\left(\frac{1}{\varphi(t)^{n-1}}\right)=\infty.$$

This is valid for this function $K$. By the definitions of $H^{-1}$ and $\psi$ we obtain that

$$\lim_{t\to 0^+} t^n K\left(\frac{1}{\varphi(t)^{n-1}}\right)=\lim_{s\to\infty}\frac{1}{s}K\left(\frac{1}{\varphi\left(s^{-\frac{1}{n}}\right)^{n-1}}\right)=\lim_{s\to\infty}\frac{K\left(H^{-1}(s)\right)}{H\left(H^{-1}(s)\right)}=\infty,$$

and thus there does not exists a constant $c$ such that

$$\|u-u_D\|_{L^K(D)}\le c\|\nabla u\|_{L^1(D)},$$

for every $u\in L^1_p(D)$.

4.12. **Theorem** (Unbounded domain, $1\le p<n$). *Assume that the function $\varphi$ satisfies the conditions* (1)– (5)*,* $C_\varphi=1$ *in the condition* (4)*, and there exists $\alpha\in[1,n/(n-1))$ such that $t^\alpha/\varphi(t)$ is increasing for $t>0$. Let the function $\psi$ be defined as in* (2.1)*. Let $D$ in $\mathbb{R}^n$, $n\ge 2$, be an unbounded domain that satisfies the following conditions:*

(a) $D=\cup_{i=1}^{\infty}D_i$*, where* $|D_1|>0$*;*
(b) $\overline{D}_i\subset D_{i+1}$ *for each* $i$*;*
(c) *each $D_i$ is a bounded $\varphi$-cigar John domain with a constant $c_J$.*

*Let $1\le p<n$. Then, there exists an $N$-function $H$, that satisfies $\Delta_2$-condition and*

$$H^{-1}(t)\approx\frac{t^{\frac{1}{p}-1}}{\psi\left(t^{-\frac{1}{n}}\right)^{n-1}}\ \textit{for all}\ t>0\,,$$

*and there exits a constant $C$ such that the inequality*

$$\inf_{b\in\mathbb{R}}\|u-b\|_{L^H(D)}\le C\|\nabla u\|_{L^p(D)},$$

*holds for every $u\in L^1_p(D)$. Here the constant $C$ depends on $n$, $p$, $C_H^{\Delta_2}$, $C_\varphi^{\Delta_2}$ and $c_J$ only.*

The proof follows the proof of [14, Theorem 4.1].

*Proof.* By Theorems 4.1 and 4.3 there exits a constant $C$ such that the inequality

$$\|u-u_{D_i}\|_{L^H(D_i)}\le C\|\nabla u\|_{L^p(D_i)} \tag{4.13}$$

holds for each $D_i$ and all $u \in L^1_p(D)$. The constant $C$ does not blow up when the diameter of $D_i$ tends to infinity. In the case $1 < p < n$ this is clear. In the case $p = 1$, we refer to the discussion after (4.7) in the proof of Theorem 4.3. The constant depends on $D_1$ but this does not cause a problem.

When $\|\nabla u\|_{L^p(D)} \le 1$ inequality (4.13) yields that there exists a constant $C < \infty$ such that the inequality

$$\int_{D_i} H(|u(x) - u_{D_i}|)\,dx \le C,$$

holds; here the constant $C$ is independent of $i$.

Let us write

$$u_i = \fint_{D_i} u(x)\,dx = \frac{1}{|D_i|}\int_{D_i} u(x)\,dx.$$

The triangle inequality yields that

$$|u_i| \le \fint_{D_1} |u(x) - u_i|\,dx + \fint_{D_1} |u(x)|\,dx.$$

Since $D_i$ satisfies inequality (4.13), we have $u \in L^H(D_1) \subset L^1(D_1)$ and thus the second term is finite. Again, by inequality (4.13) we obtain that

$$\begin{aligned}\fint_{D_1} |u(x) - u_i|\,dx &\le \frac{C\|1\|_{L^{H^*}(D_1)}}{|D_1|}\|u - u_{D_i}\|_{L^H(D_1)}\\ &\le \frac{C\|1\|_{L^{H^*}(D_1)}}{|D_1|}\|u - u_{D_i}\|_{L^H(D_i)} \le \frac{C\|1\|_{L^{H^*}(D_1)}}{|D_1|}\|\nabla u\|_{L^p(D_i)}\\ &\le \frac{C\|1\|_{L^{H^*}(D_1)}}{|D_1|}\|\nabla u\|_{L^p(D)} < \infty.\end{aligned}$$

Thus the real number sequence $(u_i)$ is bounded and hence there exists a convergent subsequence $(u_{i_j})$ and $b \in \mathbb{R}$ such that $u_{i_j} \to b$.

Since $H$ is continuous,

$$\lim_{j\to\infty} \chi_{D_{i_j}} H(|u(x) - u_{i_j}|) = \chi_D H(|u(x) - b|).$$

Fatou's lemma and the modular form of (4.13) yield that

$$\begin{aligned}\int_D H(|u(x) - b|)\,dx &\le \liminf_{j\to\infty} \int_D \chi_{D_{i_j}} H(|u(x) - u_{i_j}|)\,dx\\ &= \liminf_{j\to\infty} \int_{D_{i_j}} H(|u(x) - u_{i_j}|) \le \liminf_{j\to\infty} C = C\end{aligned}$$

for every $u \in L^1_{\mathrm{loc}}(D)$ with $\|\nabla u\|_{L^p(D)} \le 1$. This yields that there exists a constant $C$ such that the inequality

$$\|u - b\|_{L^H(D)} \le C$$

holds for every $u \in L^1_p(D)$ with $\|\nabla u\|_{L^p(D)} \le 1$. The claim follows by applying this inequality to the function $u/\|\nabla u\|_{L^p(D)}$. □

4.14. **Example.** Let the function $\varphi$ be defined as in Theorem 4.12. The following unbounded domains satisfy the assumptions of Theorem 4.12.

(a) $\{(x', x_n) \in \mathbb{R}^n : x_n \ge 0 \quad \text{and} \quad |x'| < \psi(x_n)\}$.

(b) $\mathbb{R}^2 \setminus \big(\{(x, \varphi(x)) \in \mathbb{R}^2 : 0 \le x \le 1\} \cup \{(x, -\varphi(x)) \in \mathbb{R}^2 : 0 \le x \le 1\}\big)$.

## 5. ON POINCARÉ INEQUALITIES

As a special case we recover results for Poincaré domains. We recall that a bounded domain $D$ is called a $(q, p)$-Poincaré domain, where $q, p \in [1, \infty)$, if there is a constant $C < \infty$ such that the inequality

$$\|u - u_D\|_{L^q(D)} \le C\|\nabla u\|_{L^p(D)} \tag{5.1}$$

holds for all $u \in W^{1,p}(D)$. Inequality (5.1) is the $(q, p)$-Poincaré inequality. We note that for a bounded domain $D$ inequality (5.1) holds if and only the inequality

$$\inf_{b\in\mathbb{R}} \|u - b\|_{L^q(D)} \le C_1\|\nabla u\|_{L^p(D)}$$

holds, the constants $C$ and $C_1$ depend on each other and $|D|$ only. Let us recall results for bounded $\varphi$-John domains in the case $\varphi(t) = t^s$, for a fixed $s \ge 1$. A bounded $t^s$-John domain is usually called $s$-John domain. A bounded $s$-John domain is a $(p, p)$-Poincaré domain whenever $s \in \left[1, \frac{n+p-1}{n-1}\right)$, [21, Theorem 10]. So, a bounded $s$-John domain is a $(p, p)$-Poincaré domain for all $p \ge 1$ if $s \in \left[1, \frac{n}{n-1}\right)$. A bounded $s$-John domain is a $(1, p)$-Poincaré domain if $s \in \left(1, \frac{p(n-\lambda+1)+\lambda-1}{n-1}\right)$, where $\lambda \in [n-1, n]$ is the Minkowski dimension of the boundary of the domain, [12, Theorem 1.3]. A bounded $s$-John domain is a $\left(\frac{np}{s(n-1)-p+1}, p\right)$-Poincaré domain for every $1 \le p < s(n-1) + 1$ if $s \in \left[1, \frac{n}{n-1}\right]$;[7, Corollaries 5 and 6], [17, Theorem 2.3]. The exponent $\frac{np}{s(n-1)-p+1}$ is the best possible in the class of bounded $s$-John domains, we refer to [7, p. 442]. Our Theorems 4.1 and 4.3 with $\varphi(t) = t^s$ give that a bounded $s$-John domain is a $\left(\frac{np}{n-np+sp(n-1)}, p\right)$-Poincaré domain if $1 \le p < n$ and $s \in \left[1, \frac{n}{n-1}\right)$. Thus our result is optimal in the case $p = 1$. On the other hand, our method does not cover the case $s = \frac{n}{n-1}$. Note that our proof totally differs from the previous proofs in [21, Theorem 10], [7, Corollaries 5 and 6], and [17, Theorem 2.3].

## 6. LEBESGUE SPACE CANNOT BE A TARGET SPACE

In this section we give an example which shows that for certain unbounded $\varphi$-cigar John domains the target space cannot be a Lebesgue space. The idea is that at near the infinity the target space should be $L^{np/(n-p)}$ but local structure of the domain may not allow so good integrability. We assume a priori that the function $\varphi$ has the properties (1)–(5). Later on we give extra conditions to the function $\varphi$.

We construct a mushrooms-type domain. Let $(r_m)$ be a decreasing sequence of positive real numbers converging to zero. Let $Q_m$, $m = 1, 2, \ldots$, be a closed cube in $\mathbb{R}^n$ with side length $2r_m$. Let $P_m$, $m = 1, 2, \ldots$, be a closed rectangle in $\mathbb{R}^n$ which has side length $r_m$ for one side and $2\varphi(r_m)$ for the remaining $n-1$ sides. Let $Q$ be the first quarter of the space i.e. all coordinates of the points in $Q$ are positive. We attach $Q_m$ and $P_m$ together creating 'mushrooms' which we then attach, as pairwise disjoint sets, to the side $\{(0, x_2, \ldots, x_n) : x_2, \ldots, x_n > 0\}$ of $Q$ so that the distance from the mushroom to the origin is at least 1 and at most 4, see Figure 3. We assume that a priori the function $\varphi$ has the properties (1)–(5), but we have to assume here also that $\varphi(r_m) \le r_m$. We need copies of the mushrooms. By an isometric mapping we transform these mushrooms onto the side $\{(x_1, 0, \ldots, x_n) : x_1, x_3, \ldots, x_n > 0\}$ of $Q$ and denote them by $Q_m^*$ and $P_m^*$. So again the distance from the mushroom to the origin is at least 1 and at most 4. We define

$$G = \operatorname{int}\left(Q \cup \bigcup_{m=1}^{\infty}\left(Q_m \cup P_m \cup Q_m^* \cup P_m^*\right)\right). \tag{6.1}$$

See Figure 3. We omit a short calculation which shows that $G$ is a $\varphi$-cigar John domain.



Figure 3. Unbounded $\varphi$-cigar John domain.

Let us define a sequence of piecewise linear continuous functions $(u_k)_{k=1}^{\infty}$ by setting

$$u_k(x) := \begin{cases} F(r_k) & \text{in } Q_k, \\ -F(r_k) & \text{in } Q_k^*, \\ 0 & \text{in} Q_0, \end{cases}$$

where the function $F$ will be given in (6.2). Then the integral average of $u_k$ over $G$ is zero for each $k$.

The gradient of $u_k$ differs from zero in $P_m \cup P_m^*$ only and

$$|\nabla u_k(x)| = \frac{F(r_m)}{r_m}, \text{ when } x \in P_m \cup P_m^* .$$

Note that

$$\int_G |\nabla u_k(x)|^p\,dx = 2\int_{P_m} \left(\frac{F(r_m)}{r_m}\right)^p = 2r_m\,(\varphi(r_m))^{n-1}\,\frac{F(r_m)^p}{r_m^p} .$$

We require that

$$\int_G |\nabla u_k(x)|^p\,dx = 1 .$$

Hence, we define

$$F(r_m) = \left(\frac{r_m^{p-1}}{2\varphi(r_m)^{n-1}}\right)^{1/p} . \tag{6.2}$$

Let $H$ be an $N$-function. Then,

$$\begin{aligned}\inf_{b\in\mathbb{R}} \int_G H(|u_k(x) - b|)\,dx &\geq \inf_{b\in\mathbb{R}} \Big(|Q_m| \cdot |H(F(r_m) - b)| + |Q_m^*| \cdot |H(-F(r_m) - b)|\Big)\\ &\geq r_m^n H(F(r_m)) .\end{aligned}$$

Hence, we have

$$r_m^n H(F(r_m)) = r_m^n H\bigg(\Big(\frac{r_m^{p-1}}{2\varphi(r_m^{n-1})}\Big)^{1/p}\bigg) \geq r_m^n H\bigg(\frac{1}{2}\Big(\frac{r_m^{p-1}}{\varphi(r_m^{n-1})}\Big)^{1/p}\bigg).$$

Thus, there does not exist a positive constant $C$ such that the inequality $\inf_b \|u - b\|_{L^H(G)} \leq C\|\nabla u\|_{L^p(G)}$ could hold for all $u$ from the appropriate space if

$$\lim_{t\to 0^+} t^n H\bigg(\frac{1}{2}\Big(\frac{t^{p-1}}{\varphi(t)^{n-1}}\Big)^{1/p}\bigg) = \infty.$$

Assume that $\lim_{t\to 0^+} t/\varphi(t) = \infty$. If $H(t) = t^q$, then we obtain that the inequality does not hold if

$$q \geq \frac{np}{n-p}. \tag{6.3}$$

Assume then that we have a sequence $(s_j)$ of positive numbers going to infinity. For each $s_j$ we may choose points $x(j)$ and $y(j)$ such that the balls $B(x(j), s_j)$ and $B(y(j), s_j)$ are subsets of the first quadrant and $B(x(j), 3s_j) \cap B(y(j), 3s_j) = \emptyset$. We define a sequence of piecewise linear continuous functions $(v_j)_{j=1}^\infty$ by setting

$$v_j(x) := \begin{cases} s_j^{-\frac{n-p}{p}} & \text{in } B(x_j^1, s_j),\\ -s_j^{-\frac{n-p}{p}} & \text{in } B(x_j^2, s_j),\\ 0 & \text{in } G \setminus \Big(B(x_j^1, 2s_j) \cup B(x_j^2, 2s_j)\Big) . \end{cases}$$

Now we have

$$\int_G |\nabla u_j|^p\,dx \le C s_j^n \left| \frac{s_j^{-\frac{n-p}{p}}}{s_j} \right|^p \le C$$

for some constant $C$. On the other hand, for any $b \in \mathbb{R}$

$$\begin{aligned}\int_G H(|u_j(x) - b|)\,dx &\ge C s_j^n H(|s_j^{-\frac{n-p}{p}} - b|) + C s_j^n H(| - s_j^{-\frac{n-p}{p}} - b|)\\ &\ge C s_j^n H(|s_j^{-\frac{n-p}{p}}|).\end{aligned}$$

Thus, there does not exist a positive constant $C_1$ such that the inequality $\inf_b \|u - b\|_{L^H(G)} \le C_1 \|\nabla u\|_{L^p(G)}$ could hold for all $u$ from the appropriate space if

$$\lim_{s\to\infty} s^n H(s^{-\frac{n-p}{p}}) = \lim_{s\to\infty} s^{\frac{pn}{n-p}} H\left(\frac{1}{s}\right) = \infty.$$

By choosing $H(t) = t^q$, we obtain that the inequality does not hold if

$$q < \frac{np}{n-p}. \tag{6.4}$$

If $\lim_{t\to 0^+} t/\varphi(t) = \infty$ and if there were an embedding with the Lebesgue space $L^q$ as a target space, then by (6.3) we would have $q < \frac{np}{n-p}$ and by (6.4) we would have $q \ge \frac{np}{n-p}$. Thus the target space cannot be a Lebesgue space. The target space can be $L^q$ only if $\lim_{t\to 0^+} t/\varphi(t) < \infty$. And in this case $q = \frac{np}{n-p}$. Note that the limit $\lim_{t\to 0^+} t/\varphi(t)$ exists since $\varphi$ is increasing and $\varphi \ge 0$. If $\lim_{t\to 0^+} t/\varphi(t) = m > 0$, then there exists $t_0 > 0$ such that $\frac{1}{2} m\varphi(t) \le t \le 2m\varphi(t)$.

Thus, we have proved the following theorems.

6.5. **Theorem.** *Let $\varphi$ satisfy (1)–(5), and assume that* $\lim_{t\to 0^+} t/\varphi(t) = \infty$*. Let $G$ be the unbounded $\varphi$-cigar John domain constructed in* (6.1)*. Let $1 \le p < n$. Then there do not exist numbers $q \in \mathbb{R}$ and $C \in \mathbb{R}$ such that the inequality*

$$\inf_{b\in\mathbb{R}} \|u - b\|_{L^q(G)} \le C \|\nabla u\|_{L^p(G)}$$

*could hold for all $u \in L^1_p(G)$.*

6.6. **Theorem.** *Let $\varphi$ satisfy (1)–(5), and assume that* $\lim_{t\to 0^+} t/\varphi(t) = m < \infty$*. Let $G$ be the unbounded $\varphi$-cigar John domain constructed in* (6.1)*. Assume that there exist numbers $q \in \mathbb{R}$ and $C \in \mathbb{R}$ such that the inequality*

$$\inf_{b\in\mathbb{R}} \|u - b\|_{L^q(G)} \le C \|\nabla u\|_{L^p(G)}$$

*holds for all $u \in L^1_p(G)$. Then $q = \frac{np}{n-p}$ and there exists $t_0 > 0$ such that $\varphi(t) \approx t$ for all $t \in (0, t_0]$.*

## REFERENCES

[1] A. Cianchi, *A sharp embedding theorem for Orlicz-Sobolev spaces*, Indiana Univ. Math. J. **45** (1996), 39–65.

[2] A. Cianchi, *Strong and weak type inequalities for some classical operators in Orlicz spaces*, J. London Math. Soc. (2) **60** (1999), 187–202.

[3] A. Cianchi and B. Stroffolini, *An extension of Hedberg's convolution inequality and applications*, J. Math. Anal. Appl. **227** (1998), 166–186.

[4] L. Diening, P. Harjulehto, P. Hästö, and M. Růžička, *Lebesgue and Sobolev spaces with variable exponents*, Lecture Notes in Mathematics, 2017. Springer-Verlag, Heidelberg, 2011.

[5] D. Gilbarg and N. S. Trudinger, *Elliptic Partial Differential Equations of Second Order*. Reprint of the 1998 edition. Classics in Mathematics. Springer-Verlag, Berlin, 2001.

[6] P. Hajłasz, *Sobolev inequalities, truncation method, and John domains*, Papers on Analysis: A volume dedicated to Olli Martio on the occasion of his 60th birthday. Edited by J. Heinonen, T. Kilpeläinen, and P. Koskela, Report Univ. Jyväskylä, **83**, University of Jyväskylä, Jyväskylä, 2001, pp. 109–126.

[7] P. Hajłasz and P. Koskela, *Isoperimetric inequalities and imbedding theorems in irregular domains*, J. London Math. Soc. (2)**58** (1998), no. 2, 425–450.

[8] P. Hajłasz and P. Koskela, *Sobolev met Poincaré*, Mem. Amer. Math. Soc. **145** (2000), no. 688, x+101 pp.

[9] P. Harjulehto and R. Hurri-Syrjänen, *An embedding into an Orlicz space for $L^1_1$-functions from irregular domains*, to appear in Contemp. Math. 'Complex Analysis and Dynamical Systems VI', 2015.

[10] P. Harjulehto and R. Hurri-Syrjänen, *Pointwise estimates to the modified Riesz potential*, arXiv:1406:1358.

[11] P. Harjulehto, R. Hurri-Syrjänen, and J. Kapulainen, *An embedding into an Orlicz space for irregular John domains*, F. W. Gehring Memorial Volume, Comput. Methods Funct. Theory **14** (2014), 257-277.

[12] P. Harjulehto, R. Hurri-Syrjänen, and A. V. Vähäkangas, *On the $(1,p)$-Poincaré inequality*, Illinois J. Math. **56** (2012), 905–930.

[13] L. I. Hedberg, *On certain convolution inequalities*, Proc. Amer. Math. Soc. **36** (1972), 505–510.

[14] R. Hurri-Syrjänen, *Unbounded Poincaré domains*, Ann. Acad. Sci. Fenn. A. I. Math. **17** (1992), 409–423.

[15] P. Hästö, *The maximal operator on Musielak-Orlicz spaces*, preprint, 8/2014.

[16] F. John, *Rotation and strain*, Comm. Pure Appl. Math.**14** (1961), 391–413.

[17] T. Kilpeläinen and J. Maly, *Sobolev inequalities on sets with irregular boundaries*, Z. Anal. Angew. **19** (2000), 369–380.

[18] O. Martio and J. Sarvas, *Injectivity theorems in plane and space*, Ann. Acad. Sci. Fenn. Ser. A. I. Math. **4** (1979), no. 2, 383–401.

[19] R. Näkki and J. Väisälä, *John disks*, Exposition. Math. **9** (1991), 3–43.

[20] Yu. G. Reshetnyak, *Integral representations of differentiable functions in domains with nonsmooth boundary* (Russian), *Sibirsk. Mat. Zh* **21**(1980), 108–116; translation in *Sib. Math. J.* **21** (1981), 833–839.

[21] W. Smith and D. A. Stegenga, *Hölder domains and Poincaré domains*, Trans. Amer. Math. Soc. **319** (1990), no. 1, 67–100.

[22] J. Väisälä, *Quasiconformal maps of cylindrical domains*, Acta Math. **162** (1989), 201–225.

(Petteri Harjulehto) DEPARTMENT OF MATHEMATICS AND STATISTICS, FI-20014 UNIVERSITY OF TURKU, FINLAND
*E-mail address*: `petteri.harjulehto@utu.fi`

(Ritva Hurri-Syrjänen) DEPARTMENT OF MATHEMATICS AND STATISTICS, FI-00014 UNIVERSITY OF HELSINKI, FINLAND
*E-mail address*: `ritva.hurri-syrjanen@helsinki.fi`